

OPTICAL TO X-RAYS SUPERNOVAE LIGHT CURVES FOLLOWING SHOCK BREAKOUT THROUGH A THICK WIND

GILAD SVIRSKI¹, EHUD NAKAR¹ AND RE'EM SARI²

1. Raymond and Beverly Sackler School of Physics & Astronomy, Tel Aviv University, Tel Aviv 69978, Israel
2. Racah Institute for Physics, The Hebrew University, Jerusalem 91904, Israel

Draft version February 17, 2012

Abstract

Recent supernovae (SNe) detections have motivated renewed interest in SN shock breakouts from stars surrounded by thick winds, including predictions of observable hard X-rays. Wind breakouts on timescales of a day or longer are currently the most probable for detection. Here we study the signal that follows such events, assuming a wind density profile $\propto r^{-2}$, starting from the breakout of the radiation mediated shock and tracing the evolution of the collisionless shock which forms afterwards. The emission contains two spectral components - soft (optical/UV) and hard (X-rays and possibly soft γ -rays). We find that during the breakout, the soft component temperature can vary significantly from one event to another ($10^4 - 10^6$ K), where events with longer breakout time, t_{bo} , are generally softer. The hard component is always a minute fraction, $\sim 10^{-4}$, of the breakout emission, and its fraction of the total luminosity rises quickly afterwards, gaining dominance at $\sim 10 - 50 t_{bo}$. The spectral evolution of the soft and hard components, as well as the prospects for detection of X-rays, depend mostly on the breakout time. In early breakouts ($t_{bo} \lesssim 20$ d for typical parameters) both components become harder after the breakout, and the hard component becomes dominant while the luminosity is still comparable to the breakout luminosity. In late breakouts ($t_{bo} \gtrsim 80$ d for typical parameters) the soft component becomes softer with time and the hard component becomes dominant only after the luminosity has dropped significantly. In terms of prospects for X-ray and soft gamma-ray detections, it is best to observe 100-500 days after explosions with breakout timescales between a week and a month.

1. INTRODUCTION

The breakout of a supernova (SN) shock through the stellar surface has been an active field of analytic and numerical research for several decades (e.g. Colgate 1974, Weaver 1976, Falk 1978, Klein & Chevalier 1978, Imshennik et al. 1981, Ensman & Burrows 1992, Matzner & McKee 1999). Recent advancement of observational facilities led to the discovery of several shock breakout candidates (e.g., Campana et al. 2006; Soderberg et al. 2008; Gezari et al. 2008; Schawinski et al. 2008; Modjaz et al. 2009; Ofek et al. 2010; Gezari et al. 2010; Arcavi et al. 2011) which motivated a revisiting of the theory of shock breakout (e.g., Chevalier & Fransson 2008; Katz et al. 2010; Piro et al. 2010; Nakar & Sari 2010; Katz et al. 2011a; Rabinak & Waxman 2011; Couch et al. 2011; Nakar & Sari 2011; Chevalier & Irwin 2011; Balberg & Loeb 2011; Moriya & Tominaga 2011).

The main observational challenge posed by shock breakouts from stellar surfaces is their short duration (no more than an hour even in the case of a red supergiant). However, in some cases massive mass loss prior to the SN explosion can extend the breakout signal, facilitating its detection. Ofek et al. (2010) and Chevalier & Irwin (2011) show that a breakout through a thick wind may occasionally extend long enough to account for the complete SN luminosity light curve, and suggest that at least some type II_n SNe are in fact powered by such breakouts.

When a star goes through a SN explosion, a radiation mediated shock (RMS) is driven through its envelope, accelerating through the sharp density drop near the stel-

lar edge (Sakurai 1960). If the star is not surrounded by a thick wind then the shock breaks out of the stellar surface, producing an intense short pulse of UV-X-ray photons. If, on the other hand, the star is surrounded by a wind with an optical depth between the stellar edge and the observer $> c/v$, where c is the speed of light and v is the shock speed, then the shock continues into the wind without releasing photons to the observer. At this point the shock starts decelerating and a reverse-forward shock structure is formed. The reverse shock is driven into the SN ejecta, which is characterized by a sharp rise in the amount of energy that is carried by slower moving material. The reverse shock is driving an increasing amount of energy into the shocked region, which keeps accumulating until the wind optical depth drops to c/v . Beyond this point the wind cannot contain anymore the radiation mediated forward shock (the shock width becomes $\sim R$), and the shock is breaking out of the wind, releasing all its energy as an intense pulse. This pulse can contain much more energy than a wind-less breakout, over timescales that are much longer. The general properties of a wind breakout were recently discussed by Ofek et al. (2010) and Chevalier & Irwin (2011), who find the pulse luminosity and temperature assuming thermal equilibrium, and by Balberg & Loeb (2011) who also show how the progenitor mass and explosion energy can be constrained by the pulse observational properties.

If the wind density does not fall abruptly, the energy behind the shock keeps growing after breakout, and the characteristic evolution of the resultant signal supplies an important observational probe for the interaction of

the forward shock with the wind. Katz et al. (2011b) showed that if the breakout of the RMS occurs within the wind, i.e. the matter consisting the breakout layer is spread over a scale comparable to the breakout radius rather than being concentrated within a thin layer, then a collisionless shock is bound to develop and replace the RMS in accelerating the matter ahead. This collisionless shock heats the immediate post-shock electrons to $k_B T_e \sim 60$ keV (Katz et al. 2011b), where T_e is the electron temperature and k_B is the Boltzmann constant. The cooling processes of these electrons affect the observed temperature (and possibly the luminosity evolution), which is further influenced by the interplay between photons and electrons along the photons diffusion path to the observer.

Here we examine the evolution of the luminosity and the observed temperature at and following a wind shock breakout. We consider only breakouts through a wind that is dense enough to produce a breakout pulse lasting days or longer, which are easier to detect. We also restrict the discussion here to shocks that are cooling fast during the entire evolution, i.e., the shock heated plasma cool within a dynamical time. We show that this condition is satisfied for regular SNe parameters. We pay attention to deviation from thermal equilibrium and to hydrodynamic, diffusion and cooling timescales involved, and show that the breakout timescale is the most dominant parameter in determining the post-breakout evolution. We map the different types of post breakout behaviors and describe their light curves and spectra evolution.

As this work was near completion, Chevalier & Irwin (2012) submitted a paper, presenting some of the conclusions derived below in the context of explaining the low X-ray luminosity of SN 2006gy, observed by *Chandra* at an age of 3-4 months. The low luminosity is explained as the result of three factors: two, which are discussed in details below, are the domination of inverse Compton over free-free emission in cooling the hot post-shock electrons, and the energy degradation of the hard free-free emitted photons over the diffusion path. The third is photoabsorption of soft X-rays by partially ionized pre-shock gas. This latter process is not considered in this work.

2. HYDRODYNAMIC EVOLUTION AND CHARACTERISTIC SCALES

We consider the interaction of SN ejecta with a dense stellar wind that follows a standard profile $\rho_w = Dr^{-2}$, where D is a constant and r is the radius. This wind profile implies $\tau \propto r^{-1}$, where τ is the optical depth of the wind at the location of the shock. It also implies a photon diffusion time scale to the observer, that is independent of the radius (up to a logarithmic factor), $t_{diff} \sim \kappa D/c$, where κ is the cross section per unit of mass (throughout the paper we assume that the gas which dominates the optical depth is fully ionized, and take $\kappa = 0.34 \text{ cm}^2 \text{ g}^{-1}$).

During the interaction, a forward shock is driven into the wind while a reverse shock is driven into the ejecta. Chevalier (1982) derives a self similar solution of the forward-reverse shock structure, assuming that both the ejecta and the wind density profiles are power-laws:

$$r(t) \propto t^{\frac{n-3}{n-2}} \quad (1)$$

for $\rho_w \propto r^{-2}$ and $\rho_{ej} \propto v^{-n}$. The ejecta density profile

through which the reverse shock is propagating is determined by the SN shock that crosses the stellar envelope. The density profile of the fastest moving ejecta can be approximated by a power-law with an index¹ $n \approx 10-12$ for all progenitor types. The ejecta density profile flattens towards the slower layers which contain most of the ejecta mass and energy, (e.g. Matzner & McKee 1999). As the reverse shock approaches these layers the evolution is no longer strictly self-similar, but it can be approximated by the self-similar solution using $n = 7$ (e.g., Chevalier & Irwin 2011, 2012). Given that during the reverse-forward shock interaction phase $n \gtrsim 7$ we hereafter approximate $\frac{n-3}{n-2} \approx 1$ during this phase. Finally, once the reverse shock ends crossing the ejecta only the forward shock remains, entering a Sedov-Taylor phase if the shock is adiabatic or a snowplow phase if it is radiative.

The density power-law indices of the ejecta and the wind are set by the pre-explosion stellar evolution and are not expected to vary much between various SN progenitors. The difference in the hydrodynamical evolution during the forward-reverse shock phase between various SNe is set by the two normalization factors of these density profiles. These could be determined by two observables such as the duration of the breakout pulse, t_{bo} , which is set by the wind density only, $t_{bo} \sim \kappa D/c$ (Ofek et al. 2010; Chevalier & Irwin 2011; Balberg & Loeb 2011), and the breakout luminosity L_{bo} . Below we present the evolution during the interaction phase using t_{bo} and the shock breakout velocity v_{bo} , (instead of L_{bo}) which is related to the observables via $v_{bo} = (\frac{\kappa}{4\pi c} L_{bo} t_{bo}^{-1})^{1/3}$ (Balberg & Loeb 2011). The evolution after the interaction phase ends depends on three parameters and we present it using t_{bo} , the total explosion energy, E , and ejecta mass, M_{ej} .

The breakout from the wind takes place once $\tau_{bo} \approx c/v_{bo}$, where τ_{bo} is the wind optical depth to the observer at the breakout radius. This sets the mass swept by the shock at the time of the breakout:

$$m_{bo} \approx \frac{4\pi c}{\kappa} v_{bo} t_{bo}^2 \approx 5 \times 10^{-3} M_{\odot} v_{bo,9} t_{bo,d}^2, \quad (2)$$

where $v_{bo,9} = \frac{v_{bo}}{10^9 \text{ cm/s}}$ and $t_{bo,d} = \frac{t_{bo}}{1 \text{ day}}$. During the ejecta-wind interaction the wind mass collected by the forward shock is comparable to the ejecta mass collected by the reverse shock. Therefore, using the velocity profile of an ejecta released by a SN explosion (e.g., from Nakar & Sari 2010; hereafter NS10), and requiring that a wind mass collected by the forward shock until the breakout, m_{bo} , is accelerated to v_{bo} by the explosion, implies:

$$v_{bo} \approx 10^9 \text{ cm/s } M_{10}^{-0.35} E_{51}^{0.45} t_{bo,d}^{-0.2}, \quad (3)$$

where $M_{10} = \frac{M_{ej}}{10 M_{\odot}}$ and $E_{51} = \frac{E}{10^{51} \text{ erg}}$. This equation is applicable until $m_{bo} = M_{ej}$. It implies that for typical SNe (e.g., not extremely high or low E_{51}/M_{10}) and in

¹ The mass profile for $m \ll M_{ej}$ is $m \propto v^{-\frac{\tilde{n}+1}{0.19\tilde{n}}}$ where $\tilde{n}$ (not to be confused with n) is the power-law index of the pre-explosion stellar density profile near the edge (1.5 for a fully convective envelope and 3 for a fully radiative one). This profile translates, under free expansion, to $\rho_{ej} \propto t^{-3} v^{-3-\frac{\tilde{n}+1}{0.19\tilde{n}}}$, or $n \approx 10$ for a radiative envelope and $n \approx 12$ for a convective one

the range of t_{bo} that we consider ($\gtrsim 1$ day) the breakout velocity does not vary much. Thus, we consider here only a narrow range of $v_{bo,9} = 0.3 - 1.5$. The upper limit of this range ensures that shocks that follow breakouts with timescales of days or longer remain fast cooling (i.e., the shock heated plasma radiate its energy within a dynamical time) at all times (see section 3.3). The lower velocity limit allows us to neglect recombination effects during the breakout. Interestingly, Equation 3 is independent of the progenitor radius and is very weakly sensitive to the exact density profile near the stellar edge. It is therefore applicable to red and blue supergiant as well as Wolf-Rayet progenitors. Thus, if the shock is breaking out from a thick wind, the wind density is the parameter that affects the observed radiation the most and the progenitor type cannot be identified based on the light curve alone.

An important time scale, which affects the hydrodynamic evolution, is the time at which the wind mass collected by the forward shock, m , is comparable to the total ejecta mass. At that point the reverse shock stops playing a role and the forward shock is slowing down rapidly. The deceleration rate depends on whether the shock is adiabatic, or a significant fraction of the post-shock internal energy is radiated away. If it is adiabatic then the evolution follows the Sedov-Taylor self similar solution. If, on the other hand, all the internal energy behind the shock is radiated within a dynamical time scale, then the shock enters the so called snowplow phase. We consider only fast cooling shocks, which are in the snowplow regime when $m \geq M_{ej}$ and $t > t_{bo}$.

Next we find the time of transition to snowplow, t_{SP} , in case that $t_{bo} < t_{SP}$. Equation 1 for $r(t)$ can also be applied for the snowplow phase using $n = 4$ (the Sedov-Taylor phase matches $n = 5$)². Therefore, in case that $t_{bo} < t_{SP}$ the swept mass after breakout accumulates as

$$m(t) \approx \begin{cases} m_{bo} \frac{t}{t_{bo}} & t_{bo} < t < t_{SP} \\ M_{ej} \left(\frac{t}{t_{SP}} \right)^{1/2} & t_{SP} < t \end{cases} \quad (4)$$

Since the snowplow regime begins once $m = M_{ej}$, it follows from Equation 2 that if $t_{bo} \lesssim 80 M_{10}^{0.75} E_{51}^{-0.25}$ d then $t_{bo} < t_{SP}$, where we approximate the shock velocity at the transition to the snowplow phase with $\sqrt{2E/M_{ej}}$. The transition time to snowplow phase occurs then at

$$t_{SP} = 7 \times 10^3 t_{bo,d}^{-1} M_{10}^{3/2} E_{51}^{-1/2} \text{ d} \quad (5)$$

If $m = M_{ej}$ before breakout, then the evolution follows a Sedov-Taylor expansion prior to breakout and assumes a snowplow evolution afterwards. In such case we say that $t_{SP} < t_{bo}$ and the mass behind the forward shock accumulates as

$$m(t) \approx m_{bo} \left(\frac{t}{t_{bo}} \right)^{1/2} \quad t_{SP} < t_{bo} < t \quad (6)$$

3. TEMPERATURE AND LUMINOSITY EVOLUTION

The bolometric luminosity of a fast cooling shock is $L \approx 2\pi r^2 \rho v^3 \propto v^3 \propto t^{-\frac{3}{n-2}}$. Therefore, if the breakout

² Note that using $n = 4$ in Equation 1 provides the correct evolution in the snowplow regime, although Chevalier (1982) derived it for the interaction phase only.

takes place before t_{SP} then

$$L(t) \sim \begin{cases} 5 \times 10^{43} t_{bo,d} v_{bo,9}^3 \left(\frac{t}{t_{bo}} \right)^{-0.3} \frac{\text{erg}}{\text{s}} & t \ll t_{SP} \\ 2 \times 10^{42} t_{bo,d} M_{10}^{-3/2} E_{51}^{3/2} \left(\frac{t}{t_{SP}} \right)^{-1.5} \frac{\text{erg}}{\text{s}} & t_{SP} \lesssim t \end{cases} \quad (7)$$

while if $t_{bo} > t_{SP}$ then

$$L(t) \sim 5 \times 10^{43} t_{bo,d} v_{bo,9}^3 \left(\frac{t}{t_{bo}} \right)^{-1.5} \frac{\text{erg}}{\text{s}} ; \quad t_{SP} < t_{bo} < t. \quad (8)$$

Note that once the radiation escape time becomes shorter than the dynamical time (i.e., after breakout), then as long as the shock is cooling fast, the bolometric luminosity is simply the kinetic energy flux through the shock. In contrast to regular SNe, it is independent of the gas opacity, and thus insensitive to processes such as recombination or to the gas metallicity.

Following breakout, the collisionless shock heats protons. The protons then transfer heat to electrons, which are cooled by radiation. The balance between the electrons heating and cooling rates sets the electron temperature, T_e , during the post-shock cooling process (Katz et al. 2011b). The breakout balance temperature is ~ 60 keV, independent of the shock velocity. If the protons temperature, $k_B T_p \approx \frac{3}{16} m_p v^2 \approx 200 v_9^2$ keV where m_p is the proton mass, is lower than the implied balance temperature, then $T_e \sim T_p$. Thus³

$$k_B T_{e,bo} \sim \min [200 v_{bo,9}^2, 60] \text{ keV}. \quad (9)$$

It is straight forward to show that after breakout T_e approaches T_p (if they were not similar already at breakout). Now, since we consider $v_{bo,9} = 0.3 - 1.5$, the electron and proton temperatures are not very different already during the breakout, and at $t < t_{SP}$ the proton and electron temperatures are approximately constant. At $t > t_{SP}$ the proton temperature drops, and T_e follows it. Therefore,

$$k_B T_e(t) \sim \begin{cases} 60 \text{ keV} & t < t_{SP} \\ 20 E_{51} M_{10}^{-1} \left(\frac{t}{t_{SP}} \right)^{-1} \text{ keV} & t > t_{SP} \end{cases} \quad (10)$$

While the electrons behind the collisionless shock are hot, the observed radiation at the breakout, and soon after it, is much softer (Chevalier & Irwin 2012). The reason is twofold. First, at breakout and for some time after it, the main cooling source is inverse Compton (IC) over optical-UV photons. These photons are generated by the unshocked wind that is heated by the outgoing radiation. Thus, the energy in hard photons, which are emitted by free-free emission, is suppressed by the free-free to IC cooling rate ratio. Second, even the small number of emitted hard photons do not reach the observer directly. Their energy is reduced via Compton scattering on the unshocked, lower temperature, wind electrons.

³ This equation assumes that the shock does not couple the electrons to the protons and that the electron heating is by Coulomb interactions. This is not necessarily true, in which case electrons are heated by the shock to T_p and then cool down until the radiative cooling and Coulomb heating is balanced. Since the difference between T_p and T_e is not large under any of the two scenarios their predictions for the observed signature are similar.

Thus, the observed spectrum will have two components, a soft one and a hard one. The softer component, which dominates the luminosity at early times, is produced by IC and its temperature is determined, mostly, by the deviation of the photon field in the unshocked wind from thermal equilibrium. The harder component is emitted by free-free. Its temperature is determined by the energy loss of a single photon while it diffuses out of the wind and its fraction of the total luminosity is set by the product of this energy loss and the free-free to IC cooling rate ratio. Below we calculate the evolution of the temperature of the soft component, T_{soft} , as well as the temperature and luminosity, T_{hard} and L_{hard} .

3.1. Soft component

The shock-heated electrons and protons are confined to a narrow layer behind the shock. Although they are the source of the observed luminosity, they do not necessarily set the typical photon temperature. This temperature is set during the diffusion of the photons towards the observer through the unshocked wind, as long as it is optically thick⁴. The unshocked wind is heated by the diffusing radiation, which produces photons, mostly by free-free, which share energy with the diffusing radiation. These photons also diffuse back into the hot layer behind the shock, serving as a cooling source. The soft component temperature is therefore set by the ability of the unshocked wind to produce photons over the radiation diffusion time, $t_{diff} \sim t_{bo}$. The radiation energy density at any point along the diffusion path is $\epsilon_{rad} \sim \frac{L\tau}{cr^2}$. We define $T_{BB} \equiv (\epsilon_{rad}/a)^{1/4}$, where a is the radiation constant. If photons are abundant enough and the radiation is in thermal equilibrium then the photon temperature at the shock is T_{BB} . The observer is seeing, however, a lower temperature. The reason is that thermal equilibrium is also kept at radii that are larger than the shock radius, where the radiation energy density is lower. Thus, the observed temperature is set by the radiation energy density at the largest radius at which thermal equilibrium is kept. If, in contrast, the photons at the shock are out of thermal equilibrium, then the observed temperature can be much higher than T_{BB} , and is set by the number of photons that are generated over the diffusion time. This process is the same one as in the case of a shock breakout from a stellar surface where no wind is present, and is discussed in length in NS10 and Katz et al. (2010).

Assuming that photons and electrons in the unshocked wind have the same temperature, the observed temperature can be determined using the coupling coefficient $\eta \equiv 5 \times 10^{-36} T_{BB}^{3.5} \rho^{-2} t_{diff}^{-1}$ defined in NS10, where all quantities are in c.g.s. units. Here η is the ratio of the




FIG. 1.— The observed breakout temperature as a function of the breakout velocity for $t_{bo} = 20$ d, demonstrating the strong effect of deviation from thermal equilibrium on the observed temperature. The relatively small range of relevant velocities $0.3 < v_{bo,9} < 1.5$ corresponds to a range spanning over 2.5 orders of magnitude in temperature, $10^4 \text{ K} < T_{soft,bo} < 5 \times 10^6 \text{ K}$.

photon density needed to maintain thermal equilibrium to the photons produced per unit of volume during the diffusion time, assuming that the electrons' temperature is T_{BB} . When its value at the shock radius is $\eta_s < 1$, the radiation is in thermal equilibrium and the temperature is set by the electrons at the radius where $\eta = 1$ ⁵. When $\eta_s > 1$, the observed temperature is higher than T_{BB} and is calculated as described in NS10 (with some modifications described below).

At breakout, $aT_{BB}^4/3 = (6/7)\rho_{w,bo}v_{bo}^2$, implying

$$T_{BB,bo} = \left(\frac{18c}{7\kappa a t_{bo}} \right)^{1/4} \approx 10^5 t_{bo,d}^{-1/4} \text{ K}, \quad (11)$$

and yielding

$$\eta_{bo} \approx 5v_{bo,9}^4 t_{bo,d}^{1/8}. \quad (12)$$

At any given time $\eta \propto r^{11/8}$, implying that if $\eta_{bo} \ll 1$, then the observed temperature of the breakout pulse is $T_{BB,bo}\eta_{bo}^{6/11}$, yielding

$$T_{soft,bo} \approx 20,000 \left(\frac{v_{bo}}{3000 \text{ km s}^{-1}} \right)^{2.2} t_{bo,d}^{-0.2} \text{ K} \quad \eta_{bo} < 1. \quad (13)$$

If, in contrast, $\eta_{bo} \gg 1$, then the observed temperature is much higher than T_{BB} and is given by $T_{BB,bo}\eta_{bo}^2\xi^{-2}$, where ξ accounts for the contribution of inverse Compton and its value is discussed below. Figure 1 depicts the observed breakout temperature as a function of the breakout velocity for $t_{bo} = 20$ d. It is evident that $T_{soft,bo}$ depends strongly on deviation from thermal equilibrium, rising by more than an order of magnitude when v_{bo} is increased by a factor of 2. For example, for $v_{bo,9} = 0.3$ we find $T_{soft,bo} \approx 10^4 \text{ K}$ while if $v_{bo,9} = 1.5$ then $T_{soft,bo} \approx 5 \times 10^6 \text{ K}$. The value of η that we calculate here⁶ (Equation 12) implies that the critical breakout ve-

⁴ Fast cooling dictates that the shocked plasma is cooling, forming a dense cold layer behind the hot shock heated plasma. This cold layer does not affect the observed soft component. The reason is that after the breakout the photons escape on a time that is shorter than the dynamical time, and therefore the energy density behind the shock is not dominated by radiation. Since the shock is radiative the shocked plasma is compressed to keep the pressure balance and once free-free cooling becomes dominant, its cooling rate increases with the compression and a run-away cooling decouples the plasma temperature from the rest of the system, preventing it from contributing a significant number of photons to the radiation field that cools the shock heated plasma.

⁵ We assume here that the Thompson cross section is larger than the free-free absorption cross section and that $\tau > 1$. This is true while the soft component is dominant

⁶ Note that calculating η for the configuration of a wind breakout is slightly different than in the steady state planar case discussed

locity, above which thermal equilibrium cannot be maintained, is $\sim 7000 \text{ km s}^{-1}$. However, η is calculated only to within an order of magnitude, and more importantly, the coefficient in Equation 12 may be smaller by about an order of magnitude in high metallicity environment (see discussion in NS10). Since the critical velocity for thermal equilibrium depends only weakly on that coefficient (to the power of $-1/4$), the critical velocity does not vary much between different systems. Nevertheless, since the velocity range that we consider here is exactly at the point where thermal equilibrium is marginal, we expect some of the breakouts in this range to be in equilibrium with $T_{soft,bo} \sim 10^4 \text{ K}$ and some to be out of equilibrium with $T_{soft,bo}$ in the soft X-ray range.

The evolution of T_{BB} and η_s (both measured at the shock radius) following the breakout is

$$T_{BB,s}(t) \propto \left(\frac{L\tau}{r^2}\right)^{1/4} \propto t^{-3/4}, \quad (14)$$

independent of n , and

$$\eta_s(t) \propto \frac{T_{BB,s}^{3.5}}{\rho^2} \propto T_{BB,s}^{3.5} r^4 \propto \begin{cases} t^1 & n = 12 \\ t^{0.6} & n = 7 \\ t^{-0.6} & n = 4 \end{cases}. \quad (15)$$

The evolution of the soft component while thermal equilibrium is maintained is:

$$T_{soft}(t, \eta_s < 1) \propto T_{BB,s} \eta_s^{6/11} \propto \begin{cases} t^{-0.2} & n = 12 \\ t^{-0.45} & n = 7 \\ t^{-1.1} & n = 4 \end{cases}. \quad (16)$$

Thus, while $\eta_s < 1$, the observed temperature decreases slowly at $t < t_{SP}$. It drops linearly with t during the snowplow phase for as long as $k_B T_{soft} \gtrsim 1 \text{ eV}$, beyond which recombination takes place and the temperature drop stops. When $\eta_s > 1$ (deviation from thermal equilibrium):

$$T_{soft}(t, \eta_s > 1) \propto T_{BB,s} \frac{\eta_s^2}{\xi^2} \propto \frac{1}{\xi(t)^2} \begin{cases} t^{1.2} & n = 12 \\ t^{0.4} & n = 7 \\ t^{-2} & n = 4 \end{cases}. \quad (17)$$

This equation is implicit since the factor ξ depends on T_{soft} (see below) and its evolution in time is not specified explicitly. ξ is a logarithmic factor which accounts for the contribution of inverse compton in bringing the temperature closer to thermal equilibrium (ξ always ≥ 1). It is discussed in length in NS10 and its value is ~ 1 at low temperatures ($\lesssim 10^6 \text{ K}$) and ~ 10 at high temperatures ($\gtrsim 10^7 \text{ K}$). In the scenario that we discuss here the calculation of ξ is slightly modified compared to the one described in NS10, where the softest photon that is

by Weaver (1976) and Katz et al. (2010). The reason is that in a steady state shock photons that are diffusing from the downstream back to the shock, over the shock crossing time, are produced by a mass that is larger by the shock compression ratio, 7 in this case, than the mass in the shock transition layer. In the case of a wind breakout the mass behind the shock is comparable to the mass in the shock transition layer at the time of breakout. Therefore η_{bo} is larger by a factor of ≈ 7 compared to the one that is calculated for the same shock velocity in steady state. As a result the radiation behind the wind breakout shock falls out of thermal equilibrium already when the shock exceeds 7000 km s^{-1} instead of $15,000 \text{ km s}^{-1}$ in the steady state case.

thermally coupled to the electrons is set by the requirement that its energy is doubled before it is absorbed by free-free (see their Equation 11). Here, where the optical depth of the wind can become low enough, the limiting factor may become instead the low Compton y -parameter of the wind. In our calculations below we use both requirements (high enough y -parameter and no free-free absorption) to find the softest photons that are coupled by IC to the electrons. As τ decreases the y -parameter becomes the limiting factor and the contribution of IC to coupling diminishes, so that $T_{soft}(t) \rightarrow T_{BB}\eta^2$, i.e. a very rapid temperature rise before the snowplow phase.

We note that in our calculation of T_{soft} we assume that the photons and the electrons have a single temperature at the region where T_{soft} is determined. This is always true when $\eta < 1$, but it may be broken in some cases, especially when the shock velocity is low and $\tau \ll \tau_{bo}$. If this happens then T_{soft} is higher than the value that we calculate here.

Finally, our analysis of T_{soft} is not applicable anymore when the first of the following three events takes place: (i) recombination becomes important, (ii) the optical depth drops to the point where diffusion is not a good approximation (iii) the hard component carries most of the luminosity.

3.2. Hard component

The soft component is accompanied by a harder component, which is generated by the free-free emission from the hot electrons behind the shock. At breakout, the main cooling source of these electrons is IC of soft photons, yielding a free-free to IC emissivity ratio (c.f. Chevalier & Irwin 2012):

$$\frac{\epsilon_{ff,bo}}{\epsilon_{IC,bo}} = \frac{\alpha n_e (m_e c^2)^{3/2}}{4\sqrt{k_B T_e} \epsilon_{rad}} \approx 10^{-2} v_{bo,9}^{-2}, \quad (18)$$

where n_e is the electron density, α is the fine structure constant, and $\epsilon_{rad} \approx \rho v^2/2$ at breakout. After the breakout, as long as $\tau > 1$,

$$\frac{\epsilon_{ff}}{\epsilon_{IC}}(t) \propto \frac{1}{T^{1/2} v^2 \tau} \propto \begin{cases} t & t < t_{SP} \\ t^2 & t > t_{SP} \end{cases}. \quad (19)$$

Therefore IC remains the dominant cooling source for a long time after the breakout, suppressing the emission of hard photons. But even the hard photons that are emitted at T_e are not seen directly by the observer as long as the wind optical depth is significant. Compton losses during their diffusion towards the observer limit their energy to $m_e c^2/\tau^2$ and farther reduces the luminosity of the hard component by a factor $\tau^2 T_e/m_e c^2$ (Chevalier & Irwin 2012). Therefore the observed temperature of the hard photons is

$$T_{hard}(t) = \min \left[\frac{m_e c^2}{k_B \tau^2}, T_e(t) \right], \quad (20)$$

and its luminosity is:

$$L_{hard}(t) \sim L(t) \frac{T_{hard}(t)}{T_e(t)} \min \left[1, \frac{\epsilon_{ff}}{\epsilon_{IC}}(t) \right]. \quad (21)$$

Equations 10, 18, 20 and $\tau_{bo} = c/v_{bo}$ imply that at breakout the hard component luminosity is always

$$L_{hard,bo} \sim 10^{-4} L_{bo}, \quad (22)$$

and with Equations 7 and 19 we find that at $t < t_{SP}$, while $L_{\text{hard}} \ll L$ it evolves as :

$$L_{\text{hard}}(t < t_{SP}) \propto t - t^{2.5}. \quad (23)$$

If the hard component is still subdominant during the transition to the snowplow phase, or if the breakout takes place during this phase, then its fraction out of the total luminosity rises quickly and multiplying this fraction with $L(t)$:

$$L_{\text{hard}}(t > t_{SP}) \propto t^0 - t^2, \quad (24)$$

until it becomes comparable to the bolometric luminosity.

3.3. Criterion for fast cooling

We assumed above that the shocked plasma is cooling fast (within a dynamical time). Here we find when this assumption holds. A sufficient requirement for fast cooling is that free-free emission cools the hot electrons over a dynamical time. In case the compton y -parameter drops below unity, it is also a necessary condition. The free-free cooling time at breakout is much shorter than the dynamical time for our considered range of v_{bo} :

$$\frac{t_{\text{cool},bo}^{ff}}{t_{bo}} \approx \frac{1}{25} v_{bo,9}^4 T_{e,60}^{-1/2} \quad (25)$$

Past breakout, it evolves as

$$\frac{t_{\text{cool}}^{ff}}{t} \propto v^2 n_e^{-1} T_e^{-1/2} t^{-1} \propto \begin{cases} t^{0.6} & n = 12 \\ t^{0.2} & n = 7 \\ t^{-0.5} & n = 4 \end{cases} \quad (26)$$

Thus, if the shock is cooling fast at t_{SP} then it continues to cool fast also later. Requiring $t_{\text{cool}}^{ff}(t_{SP})/t_{SP} < 1$ and taking $t_{\text{cool}}^{ff}/t \propto t^{0.6}$ ($n = 12$) all the way from breakout to t_{SP} shows that the shock enters the snowplow phase cooling fast as long as $t_{bo} \gtrsim 6 v_{bo,9}^{10/3} M_{10}^{3/4} E_{51}^{-1/4}$ d. This calculation over estimates the threshold coefficient by a factor of ~ 3 since n varies from $n \lesssim 12$ at breakout to $n \approx 4$ at t_{SP} . Therefore, in the range of breakout velocities that we consider, the shock is always fast cooling for breakout times that are days or longer.

4. LIGHT CURVES OF THE DIFFERENT REGIMES

The light curve and spectral evolution depend on three time scales. The first two are t_{bo} and t_{SP} , which affect the hydrodynamics. The third timescale, which we denote t_{hard} , marks the time at which the hard component becomes the dominant one, thus affecting the spectral evolution. For breakout times of a day or longer, the order of appearance of these three characteristic timescales depends primarily on the breakout time t_{bo} .

Since at breakout the hard component is always a small fraction of the total luminosity, $t_{bo} < t_{\text{hard}}$. Therefore, the evolution depends on the location of t_{SP} with respect to t_{bo} and t_{hard} . One regime is $t_{SP} < t_{bo} < t_{\text{hard}}$. The criterion for this regime was found in section 2. When $t_{bo} < t_{SP}$, there are two additional regimes, separated by the limiting case of $t_{SP} = t_{\text{hard}}$. In order to find the separating criterion we use Equation 21. t_{hard} is the earliest time that satisfies $L_{\text{hard}} \approx L$, namely that both $\tau^2 \approx m_e c^2 / k_B T_e$ and $\varepsilon^{ff} \approx \varepsilon^{IC}$ are satisfied. Setting $t_{SP} = t_{\text{hard}}$ the condition $\tau^2 \approx m_e c^2 / k_B T_e$,




FIG. 2.— A breakout within the snowplow regime, with $t_{bo} = 70$ d and $v_{9,bo} = 0.4$ (matching $E_{51} = 1.65$ and $m_{bo} = 10 M_{\odot}$), showing a very bright flash in the optical-UV, with a comparable rise time (not seen in the figure) and fall time. The bolometric luminosity and the temperature of the soft component (*solid lines*) decline as $t^{-1.5}$ and $t^{-1.1}$ respectively. Temperature decline stops around 6000 K, where recombination starts playing an important role (where the temperature line becomes dotted). The hard component luminosity and temperature (*dashed lines*) are strongly suppressed around the breakout (see text). $\geq$ keV photons are observed only more than a year after the explosion. The breakout time and velocity are chosen to fit the observation of SN 2006gy.

which is the last to be satisfied for typical parameters, implies $\tau \approx 5 E_{51}^{1/2} M_{10}^{-1/2}$, where we used $v(t_{SP}) \approx \sqrt{2E/M_{ej}}$ and Equation 10. Using Equation 5 and setting $\tau = 5 E_{51}^{1/2} M_{10}^{-1/2}$ one finds that $t_{SP} = t_{\text{hard}}$ when $t_{bo,d} \approx 20 M_{10}^{0.75} E_{51}^{-0.25}$. We thus obtain the following three regimes with respect to t_{SP} :

$$\begin{aligned} t_{SP} < t_{bo} & \quad t_{bo,d} \gtrsim 80 M_{10}^{0.75} E_{51}^{-0.25} \\ t_{bo} < t_{SP} < t_{\text{hard}} & \quad 20 M_{10}^{0.75} E_{51}^{-0.25} \lesssim t_{bo,d} \lesssim 80 M_{10}^{0.75} E_{51}^{-0.25} \\ t_{bo} < t_{\text{hard}} < t_{SP} & \quad 1 \lesssim t_{bo,d} < 20 M_{10}^{0.75} E_{51}^{-0.25} \end{aligned} \quad (27)$$

These regimes roughly match initial (breakout) n values of 4, 7 and 12, respectively. Below we discuss the luminosity and spectral evolution in each of these regimes.

4.1. Late breakout ($t_{SP} < t_{bo}$)

$$t_{bo,d} \gtrsim 80 M_{10}^{0.75} E_{51}^{-0.25}$$

If the progenitor went through an extreme mass loss episode just prior to the explosion, the mass in the wind can be larger than the ejecta mass. In such a scenario the breakout may take place weeks to months after the explosion, near the time that, or even after, the reverse shock is exhausted. The entire explosion energy is released then around t_{bo} . If the ejecta is several solar masses with $\sim 10^{51}$ erg, the breakout velocity is 3000 – 5000 km s $^{-1}$ and a very bright flash, with a comparable rise time and fall time, is seen in the form of optical-UV photons,

with a typical temperature of $\sim 1 - 5 \times 10^4$ K. Following breakout, the luminosity and temperature of the soft component decline as $t^{-1.5}$ and $t^{-1.1}$ respectively. The temperature decline stops around 6000 K, where recombination starts playing an important role. The breakout hard component is strongly suppressed, ($\sim 10^{-4}$ of the total luminosity) by the IC cooling dominance and the high wind optical depth. Moreover, its temperature is limited during the breakout to $\lesssim 0.5$ keV by Compton losses during the diffusion through the unshocked wind (Chevalier & Irwin 2012). The hardness and luminosity of the hard component rise following the breakout, but a significant emission above 1 keV starts only at $\sim 10t_{bo}$. Figure 2 depicts the luminosity and temperature evolution of a breakout near the snowplow phase with $t_{bo} = 70$ d and $v_{9,bo} = 0.4$ where the evolution discussed above is seen. The breakout time scale and velocity fit the observations of SN 2006gy (Ofek et al. 2007; Smith et al. 2007), which was suggested by Chevalier & Irwin (2011) to be a shock breakout through a wind. The luminosity and temperature of the dominating soft component of SN 2006gy are recovered with these t_{bo} and v_{bo} . Similarly to the conclusion of Chevalier & Irwin (2012), we find a strongly suppressed X-ray luminosity, with keV photons appearing only more than a year after the explosion (assuming that the wind is not ended at an earlier time).

Luminous hard X-ray emission is expected only if the breakout velocity is high, which requires either a low ejecta mass $\sim M_\odot$ or a very high energy explosion $\sim 10^{52}$ erg. When the breakout velocity is high and τ_{bo} is low (< 30) the hard component releases most of its energy at > 1 keV photons soon after the breakout. The luminosity of SNe in this phase is so high that X-rays may be bright enough for detection even though they carry only a small fraction of the total energy.

Note that already during the breakout the mass of the wind is comparable to the mass of the ejecta, and the mass accumulated by the shock grows as $t^{1/2}$. Therefore, if the wind mass is not much larger than the ejecta mass, the shock encounters the end of the massive wind not too long after t_{bo} .

4.2. Early breakout, ($t_{bo} < t_{hard} < t_{SP}$)

$$1 \lesssim t_{bo,d} \lesssim 20 M_{10}^{0.75} E_{51}^{-0.25}$$

If breakout occurs early the shock velocity is larger, ranging from 5,000-10,000 km s $^{-1}$ for typical SNe. The breakout emission is therefore at a marginal thermal equilibrium and a breakout temperature in the range $10^4 - 10^5$ K is expected for SNe with typical energy and ejecta mass. Figure 3 depicts the luminosity and temperature of the soft and hard components in a breakout with $t_{bo} = 10$ d and $v_{bo} = 5,000$ km s $^{-1}$. The soft component is in thermal equilibrium during breakout, $\eta_{bo} = 0.3$, resulting in $T_{soft,bo} \approx 25,000$ K. Following the breakout η_s increases roughly linearly with time, implying that it becomes harder for the radiation to maintain thermal equilibrium. As a result T_{soft} declines very slowly until day 25, when thermal equilibrium cannot be maintained anymore, and rises quickly afterwards. The luminosity decreases slowly during this time. The flux in a fixed optical/UV band is expected to drop slowly at first and then much faster. Such a breakout may explain SN PTF




FIG. 3.— Bolometric luminosity (solid line), soft component temperature (solid line), and luminosity and temperature of the hard component (dashed lines) from an early breakout of $t_{bo} = 10$ d and $v_{bo} = 5,000$ km s $^{-1}$. t_{SP} is calculated for $E_{51} = 1$ and $M_{10} = 1.25$. The temporal decay of the bolometric luminosity is shifted from $t^{-0.3}$ ($n=12$) to $t^{-0.6}$ ($n=7$) at $0.1t_{SP}$, to account for the flattening of the ejecta profile once m approaches M_{ej} . The soft component is in thermal equilibrium during breakout, $\eta_{bo} = 0.3$, resulting in $T_{soft,bo} \approx 25,000$ K. T_{soft} decreases slowly before thermal equilibrium is broken, at day 25, and rises rapidly afterwards (the soft temperature line is cut at $\tau = 3$, where diffusion approximation does not hold anymore). Such a breakout may explain SNe like PTF 09uj (Ofek et al. 2010). The hard component, very faint and soft at first, becomes brighter and harder quickly and X-rays/soft gamma-rays dominate the flux at day 200. Such events make excellent candidates for X-ray searches ~ 100 -300 d after the SN explosion.

09uj, as suggested by Ofek et al. (2010). The breakout temperature and luminosity of the case depicted in figure 3 are consistent with the observations (Ofek et al. 2010), and so does the general trend of the fading R -band flux, which becomes faster with time. The hard component is very faint at first (only $\sim 10^{-4}$ of the bolometric luminosity) and its temperature is limited to ~ 0.5 keV. But it becomes brighter and harder quickly, with hard X-rays and soft gamma-rays dominating the flux at day 200.

In the general case the breakout thermal parameter η_{bo} may be smaller or larger than the one in the case depicted in figure 3. If it is smaller then thermal equilibrium is kept for longer time, while if it is larger then the breakout temperature can be significantly larger and it starts to climb fast immediately after breakout. The hard component, as always, starts with $\sim 10^{-4}$ of the total luminosity and rises quickly to dominate the luminosity at $\sim 20t_{bo}$. Since the luminosity decays slowly during this time, these events are good candidates for X-ray searches ~ 100 -300 day after the SN explosion, when X-rays dominate the luminosity and before the snowplow phase starts. Finally, the luminosity increases with breakout time. Therefore, breakouts with larger t_{bo} (which are still in this regime) produce brighter X-ray events.

4.3. Intermediate breakout time, ($t_{bo} < t_{SP} < t_{hard}$)




FIG. 4.— An intermediate regime breakout, of $t_{bo} = 30$ d and $v_{bo} = 5,000$ km s $^{-1}$. t_{SP} is calculated for $E_{51} = 1$ and $M_{10} = 1$. The transition to the snowplow phase happens at day ≈ 230 , while the hard component gains dominance only after this transition, reaching a lower luminosity and softer X-ray temperatures compared to those of the early regime. If the breakout is early enough, so that $t_{hard} \sim t_{SP}$, such events produce the brightest X-ray luminosity among all the cases. The soft component temperature decline stops around 6000 K due to recombination (where the temperature line becomes dotted). Notations are similar to figures 2 and 3.

$$20M_{10}^{0.75}E_{51}^{-0.25} \lesssim t_{bo,d} \lesssim 80M_{10}^{0.75}E_{51}^{-0.25}$$

Figure 4 presents the luminosity and temperature of an intermediate case, where the breakout takes place before the transition to the snowplow phase, but the hard component becomes dominant only after this transition. The properties of the breakout and the emission that follows soon after it are similar to those described in the early breakout case. The difference starts at the transition to the snowplow phase, where the temperatures of both the soft and the hard components change direction and start to decrease with time. The main difference from the early breakout case is that the hard component does not dominate the luminosity at t_{SP} and by the time that it does, the luminosity is already dropping fast. If the breakout is early enough and $t_{hard} \sim t_{SP}$, then the X-ray luminosity is high. In fact these events produce the brightest X-ray luminosity among all the cases discussed in this paper. If, however, $t_{hard} \gg t_{SP}$, then the bolometric luminosity can drop significantly before X-rays dominate, resulting in a much dimmer signal.

5. SUMMARY

We examine the emission from a shock breakout through a thick wind. We restrict our exploration to breakout time scales that are days or longer, which are easier to detect, and velocities in the range 3,000-15,000 km s $^{-1}$, as expected for typical SNe. We also consider only a standard wind profile $\rho_w \propto r^{-2}$, that is extended beyond the breakout radius. Following the breakout, the internal energy in these systems is dominated by the wind material that is heated by the forward shock (Chevalier & Irwin 2011; Balberg & Loeb 2011), which

makes a transition from a radiation mediated shock to a collisionless shock at the breakout (Katz et al. 2011b). Our main conclusions are listed below.

- (i) The breakout and following emission are very luminous. In the regime that we explore, the shocked plasma is always in the fast cooling regime (see the exact condition in section 3.3), converting all the shock luminosity into radiation. Therefore, if the wind mass is comparable to, or larger than, the ejecta mass, then all the SN energy is radiated away. If the wind mass, M_w is smaller than M_{ej} then the energy radiated during the interaction is $\sim EM_w/M_{ej}$. This result supports the suggestion by Chevalier & Irwin (2011) that at least some ultra-luminous type II In SNe are such breakouts.
- (ii) The post-breakout emission is composed of two spectral components - soft and hard. The electrons behind the collisionless shock are heated to temperatures of $\sim \min[60, 200v_{bo}^2]$ keV (Katz et al. 2011b). The soft component is radiated by the IC emission of these electrons over photons produced by the heated unshocked wind, while the hard component is generated by the free-free emission of the shocked electrons.
- (iii) The soft component is in marginal thermal equilibrium during the breakout. Thus, the breakout temperature can be in the range $\sim 10^4 - 10^6$ K, where slower breakouts, $\approx 3,000$ km s $^{-1}$ will be in the lower side of this range while faster breakouts $\approx 15,000$ km s $^{-1}$ will populate the high end. In the latter case, the breakout X-ray emission of the soft component exponential tail may very well dominate over the hard component, and may possibly be detectable.
- (iv) The hard component is suppressed as long as IC is more efficient than free-free and the wind optical depth is large (Chevalier & Irwin 2012). We find that during breakout the hard component luminosity is always a fraction of $\sim 10^{-4}$ of the soft one. This result is independent of breakout time and velocity (it is also independent of the wind density profile as long as it is not terminated at the breakout radius). The fraction of the hard component out of the total luminosity rises quickly after breakout. It becomes dominant at $\sim 10 - 50t_{bo}$.

The post-breakout evolution of the two components can be divided into three regimes according to the breakout time. The behavior in each of the regimes is as follows:

Late breakout (typically $t_{bo} > 80$ d): Breakouts in the snowplow regime (i.e., when the shocked wind mass is already larger than M_{ej}) provide the brightest emission among all the regimes. All the explosion energy is released as optical-UV photons and no X-ray emission is expected during the breakout. X-rays are seen only long after the breakout in the case of regular SNe or soon after the breakout in the case of super-energetic SNe (high velocity breakout). At the peak of the hard component, around $20t_{bo}$, X-rays carry only $\sim 10^{-2}$ of the breakout energy, in the form of $\sim 1 - 10$ keV photons. Following the breakout, the temperature of the soft component drops roughly as $t^{-1.1}$ until recombination starts. The bolometric luminosity drops as $t^{-1.5}$ (also after recombination takes place), at first dominated by the soft com-

ponent, which drops quickly once the hard component becomes dominant.

Early breakout (typically $1 < t_{bo} < 20$ d): The breakout radiation is in marginal thermal equilibrium (as discussed above), and following the breakout it becomes harder to maintain thermal equilibrium. As a result, if the soft component is in thermal equilibrium at breakout, it keeps roughly a constant temperature until it falls out of thermal equilibrium, and then the temperature climbs quickly. If the soft component is out of thermal equilibrium already at breakout, then its temperature starts to rise immediately. The hard component becomes dominant before the transition to the snowplow phase, while the velocity of the shock is high and the energy behind the shock is still rising (the luminosity at this point is comparable to that of the breakout). As a result, these bursts are predicted to produce a bright hard X-rays to soft gamma-rays emission on time scales of $\sim 20t_{bo}$.

Intermediate breakout time (typically $20 \text{ d} < t_{bo} < 80 \text{ d}$): The breakout takes place before the transition to the snowplow phase, but the hard component becomes dominant only after this transition. The emission following the breakout is similar to the early breakout case until the transition to the snowplow phase, which takes place before the hard component dominates. As a result, the peak X-ray luminosity is lower than the breakout luminosity.

In our calculations we ignore the radius at which the thick wind ends. Once the wind density drops abruptly the forward shock becomes inefficient and the luminosity fades quickly (Chevalier & Irwin 2011). Since the mass of the progenitor at birth is limited, and the mass in the wind needed for an extended breakout is considerable, it is expected that the light curves we present will be terminated at some point. For example, in order for the light curve to remain bright until t_{SP} the wind mass must be

$\gtrsim M_{ej}$. Even when $M_w > M_{ej}$ the collected mass is $\propto t^{1/2}$ after t_{SP} , limiting the lifetime of the bright emission. In terms of the prospects of observing X-rays, this consideration boosts the attractiveness of earlier breakouts, since the mass that early breakout events collect by the time that the hard component dominates is lower.

We do not address the extinction of soft X-rays due to photoabsorption by partially ionized unshocked wind (Chevalier & Irwin 2012). This process may suppress softer-end photons of the hard component as well as hard photons of the soft component, if the latter is far from thermal equilibrium. If photoabsorption is important then the observed flux of soft X-rays is lower than the one that we predict.

Finally, our treatment ignores the contribution of emission from ejecta mass before it is shocked by the reverse shock. Once the Compton y -parameter of the wind drops below unity, this emission becomes similar to the emission from a typical SN without a thick wind (e.g., typical IIP, Ib and Ic SNe). The emission from the forward shock that we consider here always outshines this emission (in terms of bolometric luminosity). But, at late times, when most of the forward shock emission is in hard photons, the contribution from inner layers may dominate the IR-optical emission.

To conclude, breakouts through thick winds produce very bright SNe with a potential for detection across the entire electromagnetic spectrum. The brightest phase is the breakout, which is always dominated by optical-UV emission with a negligible X-ray emission. The X-rays and even soft gamma-rays luminosity rises after the breakout, and reaches its peak after $\sim 10 - 50 t_{bo}$.

We thank Eran Ofek and Boaz Katz for helpful discussions and comments. G.S. and E.N. were partially supported by an ISF grant (174/08) and by an ERC starting grant (GRB-SN 279369). R.S. was partially supported by ERC and IRG grants, and a Packard Fellowship.

REFERENCES

- Arcavi, I., et al. 2011, *ApJL*, 742, L18
 Balberg, S., & Loeb, A. 2011, *MNRAS*, 414, 1715
 Campana, S., et al. 2006, *Nature*, 442, 1008
 Chevalier, R. A. 1982, *ApJ*, 258, 790
 Chevalier, R. A., & Fransson, C. 2008, *ApJL*, 683, L135
 Chevalier, R. A., & Irwin, C. M. 2011, *ApJL*, 729, L6+
 —. 2012, *ArXiv e-prints*
 Colgate, S. A. 1974, *ApJ*, 187, 333
 Couch, S. M., Pooley, D., Wheeler, J. C., & Milosavljević, M. 2011, *ApJ*, 727, 104
 Ensman, L., & Burrows, A. 1992, *ApJ*, 393, 742
 Falk, S. W. 1978, *ApJL*, 225, L133
 Gezari, S., et al. 2008, *ApJL*, 683, L131
 —. 2010, *ApJL*, 720, L77
 Imshennik, V. S., Nadezhin, D. K., & Utrobin, V. P. 1981, *Ap&SS*, 78, 105
 Katz, B., Budnik, R., & Waxman, E. 2010, *ApJ*, 716, 781
 Katz, B., Sapir, N., & Waxman, E. 2011a, *ArXiv e-prints*
 —. 2011b, *ArXiv e-prints*
 Klein, R. I., & Chevalier, R. A. 1978, *ApJL*, 223, L109
 Matzner, C. D., & McKee, C. F. 1999, *ApJ*, 510, 379
 Modjaz, M., et al. 2009, *ApJ*, 702, 226
 Moriya, T. J., & Tominaga, N. 2011, *ArXiv e-prints*
 Nakar, E., & Sari, R. 2010, *ApJ*, 725, 904
 —. 2011, *ArXiv e-prints*
 Ofek, E. O., et al. 2007, *ApJL*, 659, L13
 —. 2010, *ApJ*, 724, 1396
 Piro, A. L., Chang, P., & Weinberg, N. N. 2010, *ApJ*, 708, 598
 Rabinak, I., & Waxman, E. 2011, *ApJ*, 728, 63
 Sakurai, A. 1960, *Communications on Pure and Applied Mathematics*, 13, 353
 Schawinski, K., et al. 2008, *Science*, 321, 223
 Smith, N., et al. 2007, *ApJ*, 666, 1116
 Soderberg, A. M., et al. 2008, *Nature*, 453, 469
 Weaver, T. A. 1976, *ApJS*, 32, 233